

Long Range PCR protocol for KOMP-CSD derived ESC lines

September 8th 2009

I DNA extraction from ES cells (following manufacturers suggestions)

Materials

DNeasy® Tissue Kit 250 (Qiagen 69506) PBS (50mM potassium phosphate; 150mM NaCl)

Procedure (12-well plates)

1. Wash the monolayer with cold PBS (4°C).
2. Scrape the cells with 0.5 ml of cold PBS using a rubber policeman, and transfer to 1.5ml tube on ice.
3. Add 20 µl proteinase K and 200 µl Buffer AL (without added ethanol). Mix thoroughly by vortexing, and incubate at 70°C for 10 min. It is essential that the sample and Buffer AL are mixed immediately and thoroughly by vortexing or pipetting to yield a homogeneous solution.
4. Add 200 µl ethanol (96–100%) to the sample, and mix thoroughly by vortexing. It is important that the sample and the ethanol are mixed thoroughly to yield a homogeneous solution.
5. Pipet the mixture from step 3 into the DNeasy Mini spin column placed in a 2 ml collection tube (provided). Centrifuge at $\approx 6000 \times g$ (8000 rpm) for 1 min. Discard flow-through and collection tube.
6. Place the DNeasy Mini spin column in a new 2 ml collection tube (provided), add 500 µl Buffer AW1, and centrifuge for 1 min at $\approx 6000 \times g$ (8000 rpm). Discard flow-through and collection tube.
7. Place the DNeasy Mini spin column in a new 2 ml collection tube (provided), add 500 µl Buffer AW2, and centrifuge for 3 min at 20,000 $\times g$ (14,000 rpm) to dry the DNeasy membrane. Discard flow-through and collection tube. It is important to dry the membrane of the DNeasy Mini spin column, since residual ethanol may interfere with subsequent reactions.
8. Place the DNeasy Mini spin column in a clean 1.5 ml microcentrifuge tube, and pipet 100 µl Buffer AE directly onto the DNeasy membrane. Incubate at room temperature for 1 min, and then centrifuge for 1 min at 8000 rpm to elute.
9. Store at -4° C until use or -20° C if storing more than a day.

II Long Range PCR

Universal Oligos: (5' to 3')

3' Universal (R2R): TCTATAGTCGCAGTAGGCGG (fwd from genetrapp cassette)

5' Universal (LAR3): CACAACGGGTTCTTCTGTAGTCC (rev from genetrapp cassette)

Access to clone specific vector and oligo data:

1. From KOMP.org type official gene symbol into search bar and click on correct gene symbol when list populates.
2. Under "Show All Projects" click on the 8 digit project ID "CSDXXXXX" relating to your order.
3. Use any or all 5' Gene Specific (GF) in conjunction with 5' Universal (LAR3).
4. Use any or all 3' Gene Specific (GF) in conjunction with 3' Universal (R2R).
5. Access the Genbank File by clicking on "view" under ES Cell Clones bar. This will provide sequence to use for amplicon size expectation.

KOMP Repository
Knock Out Mouse Project 1-888-KOMP-MICE

kdawson Logged In Home MyKOMP Logout Utilities Storage Projects Reports

Abca15

ATP-binding cassette, sub-family A (ABC1), member 15
MGI:2388709 (Current, syntenic, Gene)


Ensembl: [E177213](#)

Pipeline Summary: CSD: ES cells - Targeting confirmed | KOMP

Pre-pipeline Design Strategy Vector Construction ESC Electroporation Tra

CSD33366 Complete

CSD33367 ESC Targeting Confirmed



LR PCR Genotyping Primers

5' Gene Specific (GF3)	GTCCAGTCCACAAGCACCTACCTCAGC
5' Gene Specific (GF4)	GTCCAGTCCACAAGCACCTACCTC
5' Universal (LAR3)	CACAACGGGTTCTTCTGTAGTCC
3' Universal (RAF5)	CACACCTCCCCCTGAACCTGAAC
3' Universal (PIF)	ATCCGGGGTACCAGCTCGAG
3' Universal (R2R)	TCTATAGTCGCAGTAGGCGG
3' Gene Specific (GR3)	GTATTGGAGATTGGCCAGGTTTC
3' Gene Specific (GR4)	CATTGTATTGGAAGATTGGCCAGGTTTC

Materials and equipment

SequelPrep long PCR kit, Invitrogen A10498 Peltier Tetrad2 thermal cycler, BioRad. Oligos, Invitrogen

Procedure

1. Thaw and equilibrate all buffers at room temperature
2. Prepare the following master mix on ice (1X shown):

Components	Volume per rxn
Sterile H2O	13.24
10X Buffer (green tube)	2
Enhancer A (Red tube)	1
Enhancer B (Yellow tube)	1
DMSO (brown tube)	0.2
gene specific primer (10uM)	0.5
universal primer (10uM)	0.5
Enzyme (black tube)	0.36
total cocktail	19
template	1
reaction volume	20

3. Briefly vortex master mix and transfer 19 ul of to each 200ul thin walled reaction tube on ice.
4. Briefly flick DNA tube and input 1 ul of ~100 ng into reaction tube.
5. PCR with the following thermal conditions:
Note: may adjust elongation time to 1min/kb amplicon.

Temp	Duration	Repetitions
93°C	3 minutes	1x
93°C	15 seconds	8X
68°C	30 seconds -1°C/cycle	
68°C	9 minutes	30X
93°C	15 seconds	
60°C	30 seconds	
68°C	9 min + 20 sec/cycle	1x
68°C	9 minutes	
4°C	hold	

III Data Analysis

Materials and Equipment

Loading Dye (15ml glycerol; 35ml H ₂ O, 125mM each Bromophenol Blue/Xylene Cyanol).
1 Kb plus DNA Ladder, Invitrogen 10787-026.
GenePure LE Agarose, ISCBioExpress E-3120-500.
Gel Logic Imaging System 100, Kodak.

Procedure

1. Prepare a 1% Agarose gel with 0.25ug Ethidium Bromide per ml agarose. (non toxic comparable would be SYBRsafe, Invitrogen S33102).
2. Line pipette tips with Loading Dye and mix with finished PCR reaction.
3. Load 15ul of reaction into well.
4. Run gel at 120 volts for 2 hours.

5. Image under UV and adjust and store image for record.

6. Controls: non-template control, isogenic wild type, 1kb plus ladder.
(no positive control available).